RESEARCH ARTICLE

Mycorrhizal Formation of Korean Truffle Species, *Tuber koreanum* with Oak Trees

Ju-Hui Gwon, and Ahn-Heum Eom

Department of Biology Education, Korea National University of Education, Cheongju 28173, Korea

^{*}Corresponding author: eomah@knue.ac.kr

ABSTRACT

In this study, we examined the ectomycorrhiza (ECM) formed by a species of the genus *Tuber, T. koreanum*, with two indigenous Korean oak species, *Quercus acutissima* and *Q. dentata. T. koreanum* showed a high colonization rate for both oak species, with a higher mean colonization rate observed for *Q. acutissima*. The mycorrhizal characteristics of *T. koreanum* were described, revealing distinct morphological and anatomical differences between the ECM formed with the two host plants. This study is significant because it is the first to elucidate ECM formations of *T. koreanum* in indigenous oak trees. However, further studies are required to differentiate the ECM associations formed by *T. koreanum* from those of other *Tuber* species.

Keywords: Ectomycorrhiza, Inoculated seedlings, Oaks, Truffle, Tuber koreanum



OPEN ACCESS

pISSN: 0253-651X **eISSN**: 2383-5249

Kor. J. Mycol. 2023 June, 51(2): 121-125 https://doi.org/10.4489/KJM.20230013

Received: May 30, 2023 Revised: June 18, 2023 Accepted: June 19, 2023

© 2023 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access

article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http: //creativecommons.org/licenses/by-nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. **INTRODUCTION**

The genus *Tuber* (Pezizales, Ascomycota) comprises over 180 species worldwide [1,2]. These fungi form ectomycorrhizal (ECM) relationships with certain host plants, such as oak or hazel [3], leading to the production of subterranean fruiting bodies [4]. Although most ECM fungi form symbiotic relationships with a number of different host plant species [5,6], *Tuber* spp. exhibit high selectivity for specific hosts [1]. For example, *Tuber gibbosum* and *T. oregonense* establish unique symbiotic relationships with *Pseudotsuga menziesii* [2].

In Greece, the optimal host species for truffle cultivation include *Quercus coccifera*, *Q. ilex*, and *Q. pubescens* [7]. Indeed, careful selection of the host species is crucial for successful truffle cultivation. Moreover, each *Tuber* species exhibits unique morphological or anatomical characteristics when forming ECM associations, underlining the importance of understanding these characteristics for their cultivation [8,9].

In September of 2020, fruiting bodies were collected from the rhizosphere of *Q. aliena* in Gyeongju, Korea. Based on their morphological characteristics and molecular analysis, we identified *T. koreanum* as a new *Tuber* species [10]. Since then, *T. koreanum* has also been found in the rhizosphere of *Q. aliena* in both Mungyeong and Uljin.

In the present study, we sought to characterize the ECM associations formed by *T. koreanum* when inoculated into two different oak tree species commonly found in Korea, *Q. acutissima* and *Q. dentata*. *Quercus acutissima* was chosen as it is representative of the oak tree species frequently found in Korea, whereas *Q. dentata* was selected owing to its role as a host plant for other *Tuber* spp. indigenous to Korea [11,12].

MATERIALS AND METHODS

Seeds of *Q. acutissima* were collected from Danyang-gun, Korea, whereas those of *Q. dentata* were collected from Yeongju-si, Korea. The collected seeds were then de-shelled and their surfaces sterilized in 10% sodium hypochlorite (NaOCl) for 30 min. Following sterilization, each seed was placed in a plastic pot (SC10, Stuewe and Sons., Inc, Oregon, USA) containing an autoclaved soil mixture (1:1 ratio of vermiculite to perlite, sterilized for 1 h at 121°C) and incubated in a greenhouse at 25°C. Subsequently, seeds were maintained in a growth chamber under controlled conditions (8-h photoperiod per day, $55\pm5\%$ relative humidity, and $24\pm1°C$ temperature) for 8 months.

The fruiting bodies of *T. koreanum* were collected from the rhizosphere of *Q. aliena* in Gyeongju-si, Korea [10]. The *T. koreanum* ascocarps were rinsed with tap water, sterilized with 70% EtOH, and ground with sterile water using a blender (SGMF-650, Hanil, Seoul, Korea), resulting in a spore suspension. Each previously prepared seedling was inoculated with 1 mL of *T. koreanum* spore suspension $(1.0 \times 10^5 \text{ spores/mL})$. At the time of inoculation, the spore suspension was applied near the root. In total, ten inoculated seedlings were produced, comprised of two different species combinations (*Q. acutissima* × *T. koreanum* and *Q. dentata* × *T. koreanum*) with five trees each. The growth area of the inoculated seedlings was uniformly maintained, the soil was watered to retain moisture, and the pH was maintained at 8 by applying with slaked lime.

Approximately 8 months of inoculation, the ECM-associated root tips were classified based on their morphological characteristics, and representative root tips from each group were selected for molecular identification. To determine the successful formation of *T. koreanum* ECM in each inoculated seedling, DNA was extracted from the root tips using a DNeasy Plant Mini kit (Qiagen GmbH, Hilden, Germany). Polymerase chatin reaction (PCR) was conducted using ITS1F/ITS4 primers [13] and DNA sequencing was performed (SolGent Co., Ltd., Daejeon, Korea). Using NCBI BLAST (https://www.ncbi.nlm.nih.gov/), species showing the highest agreement were identified for the nucleotide sequence analysis.

Colonization rates (CRs) were assessed eight months after inoculation. From each seedling, the soil near the root was removed and washed with distilled water, and three root sections (2-3 cm each) were randomly sampled. The colonization rate was calculated by counting the total number of *T. koreanum*, ECM root parts, non-ECM root parts, and contaminated root parts from the extracted roots. A total of 100 root sections from each plant species were examined, and the results are expressed as the percentage of tips infected with *T. koreanum*. Statistical analysis were performed using SPSS (version 26, IBM Corp., NY, USA). Statistical differences between the measured data of each host plant were compared using an independent student t-test.

Eight months after inoculation, the mycorrhizal characteristics of *T. koreanum* formed on the roots of each seedling were observed. The morphological characteristics of the ECM were examined and recorded using a stereomicroscope (Olympus SZX7, Olympus, Tokyo, Japan). Sections were prepared using a frozen microtome (LEICACM1850, Leica, Heidelberg, Germany), and ECM anatomical properties were observed and recorded using an optical microscope (Axio Imager A1, Carl ZEISS, Oberkochen, Germany).

RESULTS AND DISCUSSION

Approximately 8 months after inoculation with the *T. koreanum* spore suspension, mycorrhizal formation was observed. For both host plant species, *T. koreanum* successfully colonized all five seedlings. The BLAST results showed that the internal transcribed spacer (ITS) sequences of all observed ECM associations were closely related to those of the ascoma (GenBank accession no. AB553502), with more than 98% similarity, confirming that the DNA sequence of the ECM belonged to the same phylogenetic group as the ITS sequence of the ascoma inoculated.

Upon measuring CRs in both host plant species, the mean CR for the *Q. acutissima* \times *T. koreanum* combination was 0.46±0.09, whereas that for the *Q. dentata* \times *T. koreanum* combination was 0.31±0.03. While the average CR appeared to be higher in *Q. acutissima* than that in *Q. dentata*, the difference was not statistically significant.

Eight months after inoculation, the ECM formed in *Q. acutissima* and *Q. dentata* were examined. The morphological and anatomical characteristics of the ECM formed in each host plant are presented in Fig. 1, and the corresponding measured values are listed in Table 1. In both host plant species, the unramified ends were straight, club shaped, or cylindrical with rounded ends. The mantle is characterized by an irregular pattern of interlocking hyphae, which are simple, bristle-like, awl-shaped, and non-septate.

Different mycorrhizal properties were observed in the ECM associations formed with the two host plant species. Both monopodial pinnate and pyramidal patterns were observed in that of *Q. acutissima*, whereas only the monopodial pinnate pattern was observed in that of *Q. dentata*. The ECM of *Q. acutissima* was whitish-light brown, brown, or reddish-brown in color, whereas the ECM of *Q. dentata* was either pale yellow or ochre. Furthermore, the ECM system was significantly longer in *Q. dentata* than in *Q. acutissima*. Additionally, the length and diameter of the unramified ends and cystidia were significantly greater in *Q. dentata* than in *Q. acutissima*. In contrast, mantle thickness was significantly larger in *Q. acutissima* than in *Q. dentata* (Table 1).

Most *Tuber* spp. exhibit host selectivity at the species level [14]. In fact, some species have been found to have higher affinity with certain species of oak tree than others [7,15]. When *T. koreanum* was inoculated into indigenous host plant species and the CRs were compared, a higher CR was observed in *Q. acutissima* than in *Q. dentata*. These findings suggest that *T. koreanum* may exhibit a preference for this specific host plant species. Selecting suitable indigenous host plants is important for the successful mycorrhization of *Tuber* spp. [1]. Although this study compared the CRs of two species, further studies comparing the CRs of a larger range of host plant species are warranted. This study will facilitate more effective cultivation of *T. koreanum* by helping to identify optimal host plant species.



Fig. 1. Macro-morphological and anatomical characters of *Tuber koreanum* mycorrhizae with *Quercus acutissima* (a-d) and Q. dentata (e-h). The shape of mycorrhizal root tips (a & e); Cross section of mycorrhizal root tips (b & f); Outer mantle surface structure (c & g); Separate hyphae emanating from outer mantle layer (d & h) (scale bars: a, e=1 mm; b, c, d, h=50 µm; f, g=20 µm).

Dimensions	Quercus acutissima	Quercus dentata	p-value ^b
ECMs system (mm)	4.56 ± 0.63	7.28 ± 0.39	0.040
Length of unramified ends (mm)	200.00 ± 4.85	268.50 ± 3.50	< 0.001
Diamenter of unramified ends (µm)	225.67 ± 5.49	257.17 ± 3.87	< 0.001
Thick of mantle (µm)	33.33 ± 1.82	17.10 ± 2.27	< 0.001
Length of cystidia (µm)	154.17 ± 8.80	$212.50 \pm 1 \ 0.94$	0.002
Diameter of cystidia (µm)	$1.45~\pm~0.04$	$2.23~\pm~0.05$	< 0.001

ECM: ectomycorrhiza.

 $amean \pm standard error, n=100, bmp-values were obtained from statistical analysis using independent t-test.$

In the present study, we investigated the formation and characteristics of ECM produced by *T. koreanum* when associated with either *Q. acutissima* or *Q. dentata*. For the first time, this study describes ECM formation of *T. koreanum* in association with indigenous oak trees. However, distinguishing between the ECM formed by *T. koreanum* and that formed by other *Tuber* species remains challenging, underscoring the need for further studies on the formation and characteristics of ECM.

ACKNOWLEDGMENT

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (IPET319106052HD050).

REFERENCES

- 1. Bonito GM, Trappe JM, Rawlinson P, Vilgalys R. Improved resolution of major clades within *Tuber* and taxonomy of species within the *Tuber gibbosum* complex. Mycologia 2010;102:1042-57.
- Bonito GM, Gryganskyi AP, Trappe JM, Vilgalys R. A global meta-analysis of *Tuber* ITS rDNA sequences: species diversity, host associations and long-distance dispersal. Mol Ecol 2010;19:4994-5008.
- Núñez JAD, Serrano JS, Barreal JAR, de Omeñaca González JAS. The influence of mycorrhization with *Tuber melanosporum* in the afforestation of a Mediterranean site with *Quercus ilex* and *Quercus faginea*. For Ecol Manag 2006;231:226-33.
- Urban A. Truffles and small mammals. True truffle (*Tuber* spp.) in the world. Berlin: Springer; 2016. p. 353-73.
- 5. Horton TR, Bruns TD. Multiple-host fungi are the most frequent and abundant ectomycorrhizal types in a mixed stand of Douglas fir (*Pseudotsuga menziesii*) and bishop pine (*Pinus muricata*). New Phytol 1998;139:331-9.
- Ishida TA, Nara K, Hogetsu T. Host effects on ectomycorrhizal fungal communities: insight from eight host species in mixed conifer–broadleaf forests. New Phytol 2007;174:430-40.
- Tsiaras S, Dragoslis A, Papathanasiou J. Fuzzy multiple criteria analysis for selecting the optimum tree species for truffle cultivation in Greece. In: International Workshop "Information Technology, Sustainable Development, Scientific Network and Nature Protection". Edessa: 18th Panhellenic Forestry Congress; 2017.
- 8. Kovács GM, Jakucs E. Morphological and molecular comparison of white truffle ectomycorrhizae. Mycorrhiza 2006;16:567-74.
- Guerin-Laguette A, Cummings N, Hesom-Williams N, Butler R, Wang Y. Mycorrhiza analyses in New Zealand truffières reveal frequent but variable persistence of *Tuber melanosporum* in co-existence with other truffle species. Mycorrhiza 2013;23:87-98.
- Park H, Gwon JH, Lee JC, Eom AH. Report on a new truffle species, *Tuber koreanum* sp. nov., from Korea. Mycobiology 2021;49:527-33.
- 11. Park H, Gwon JH, Lee JC, Kim HS, Oh DS, Eom AH. Report on *Tuber huidongense*, a truffle species previously unrecorded in Korea. Kor J Mycol 2020;48:505-10.
- Park H, Gwon JH, Lee JC, Kim HS, Seo GS, Eom AH. Morphological and phylogenetic characteristics of *Tuber himalayense* collected from rhizosphere of *Quercus dentata* in Korea. Kor J Mycol 2021;49:101-8.
- 13. Gardes M, Bruns TD. ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. Mol Ecol 1993;2:113-8.
- Molina R, Trappe JM. Patterns of ectomycorrhizal host specificity and potential among Pacific Northwest conifers and fungi. For Sci 1982;28:423-58.
- 15. Bonet JA, Fischer CR, Colinas C. Cultivation of black truffle to promote reforestation and land-use stability. Agron Sustain Dev 2006;26:69-76.