

RESEARCH NOTE

Erysiphe quercicola Causing Powdery Mildew on *Quercus rubra* in Korea

Young-Joon Choi¹, Jun Hyuk Park¹, In-Young Choi², Lamiya Abasova², Joon-Ho Choi², and Hyeon-Dong Shin^{2,3,*}

¹Department of Biological Science, Kunsan National University, Gunsan 54150, Korea

²Department of Agricultural Biology, Jeonbuk National University, Jeonju 54896, Korea


³Division of Environmental Science and Ecological Engineering, Korea University, Seoul 02841, Korea

*Corresponding author: hdshin@korea.ac.kr

ABSTRACT

In Korea, powdery mildew has been observed on *Quercus rubra*, commonly called the northern red oak tree. Based on its morphological and molecular phylogenetic analyses, the fungal agent was identified as *Erysiphe quercicola*, but it was distinct from the other four *Erysiphe* species, specialized to *Quercus* species. This is the first confirmed report on the identity of powdery mildew found on *Q. rubra* in Korea. This finding can help understand the host expansion of *E. quercicola* and the biodiversity of powdery mildew on *Quercus* species.

Keywords: Emerging disease, Host expansion, Morphology, Phylogeny



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2022 June, 50(2): 137-141
<https://doi.org/10.4489/KJM.20220014>

Received: March 07, 2022

Revised: June 09, 2022

Accepted: June 13, 2022

© 2022 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Quercus rubra L., commonly called the northern red oak, is native to North America and is cultivated as an ornamental tree in temperate regions. It is widely grown in parks and gardens in Korea. August 2018, powdery mildew was observed on several *Q. rubra* trees in a public garden in Seoul, South Korea. White superficial fungal colonies were present on both sides of the leaves and young shoots (Fig. 1A). Severe infections often caused malformation and necrotic discoloration of the leaves (Fig. 1B). No chasmothecia were observed until the infected leaves dropped and dried in December (Fig. 1C).

For morphological examination, anamorphic structures were detached from the powdery mildew colonies on fresh leaves of the plant using a sterile razor blade, mounted in distilled water, and examined using bright-field and differential interference contrast (DIC) light microscopy (Carl Zeiss, Oberkochen, Germany). At least 20 measurements were performed for each morphological structure. Dried herbarium samples were analyzed using the lactic acid technique [1]. Two voucher specimens were deposited at the Korea University Herbarium in Seoul, Korea, under accession numbers KUS-F30755 and F30867.

The morphological characteristics of the fresh samples were as follows: The hyphae were septate, branched, and 4–8 µm in width. Hyphal appressoria were multilobed to moderately lobed, single or in pairs, and 4–7 µm wide (Fig. 1D). Conidiophores were 3–4-celled, cylindrical, 56–98 × 7–9 µm in size, and produced conidia singly (Fig. 1E–G). Foot cells of conidiophores were cylindrical, straight, or slightly flexuous at the base and 24–46 µm long, with a basal septum at 6–12 µm away from the branching point

of the mycelium. Conidia were ellipsoid or doliiform, $25\text{--}38 \times 18\text{--}24\text{ }\mu\text{m}$ with a length/width ratio of 1.5–1.9 and devoid of distinct fibrosin bodies (Fig. 1K–P). The primary conidia were apically round and basally sub-truncated (Fig. 1H–J). Germ tubes were formed on the perihilar position of the conidia




Fig. 1. *Erysiphe quercicola* causing powdery mildew on *Quercus rubra*. A: Affected tree showing powdery mildew symptoms. B: Close-up of affected leaves. C: Fallen leaves with powdery mildew symptoms. D: Appressoria on a hypha. E–G: Conidiophores. Arrows indicate septa. H–J: Primary conidia. K–P: Conidia. Q–R: Germinating conidia.

(Fig. 1Q–R). The structure and measurements were consistent with the anamorphic features of *Erysiphe quercicola* S. Takam. & U. Braun [2,3].

For molecular analysis, genomic DNA was extracted from conidia and mycelia harvested from the infected leaves of KUS-F30755 using a MagListo 5 M plant Genomic DNA Extraction Kit (Bioneer,

Daejeon, Korea). The nucleotide sequences of the internal transcribed spacer (ITS) and 28S rDNA regions were amplified using primer pairs PM10/ITS4 and PM3/TW14, respectively, under the PCR conditions outlined by Bradshaw and Tobin [4]. The PCR products were purified using an AccuPrep PCR Purification Kit (Bioneer) and sequenced by Macrogen Inc. (Seoul, Korea). The resulting sequences were edited using the DNASTar software package (Madison, Wis., USA) version 5.05 and deposited in GenBank (Accession nos. OM899628 and OM899630). A BLASTn search was conducted to investigate sequence similarities with reference sequences in NCBI GenBank. The sequences of individual markers, including the reference sequences of *Erysiphe* species originating from *Quercus* spp. retrieved from GenBank, were aligned using MAFFT 7 [5] and concatenated in SequenceMatrix v1.7.8 [6]. A phylogenetic tree of the concatenated alignment was constructed by the minimum evolution (ME) method using MEGA X [7] with the default settings of the program, except by using the Tamura-Nei model instead of the maximum




Fig. 2. Phylogenetic tree of *Erysiphe* species from the minimum evolution analysis based on a concatenated alignment of the internal transcribed spacer (ITS) and large subunit (LSU) of rDNA sequences. Bootstrap values higher than 60% are shown above the branches (1,000 replicates). The green box covers the sequences of *Erysiphe quercicola* and Korean isolates obtained from *Quercus rubra* are displayed in bold. The scale bar indicates the number of nucleotide substitutions per site.

composite likelihood model. The robustness of the phylogenetic tree was evaluated using 1,000 bootstrap (BS) replicates.

Based on a BLASTn search of the NCBI GenBank, the Korean specimen showed 100% sequence similarity (668/668 bp) with the reference sequences of *E. quercicola* (AB193607, AB237808, AB292693, LC270856, MH333267) for ITS and 99.9-100% (997 or 998/998 bp) similarity with those of *E. quercicola* (AB292692, MN1659866-7, LC128423-5) for LSU sequences. ME analyses of a concatenated alignment of ITS and LSU sequences were performed to infer the phylogenetic relationships between the Korean specimen and previously identified *Erysiphe* species parasitic on *Quercus* spp. The phylogenetic tree (Fig. 2) revealed that the Korean specimens grouped with the reference sequences of *E. quercicola*, parasitic on *Q. crispula*, *Q. phillyraeoides*, and *Q. serrata*, with the maximum bootstrapping value, but were distantly related to the other four species of *Erysiphe* parasitic on *Quercus* species, namely, *E. alphitoides*, *E. epigena*, *E. hypogena*, and *E. hypophylla*.

A literature review showed that two species of *Erysiphe* (including formerly *Microsphaera*) have been recorded as powdery mildews on *Q. rubra* [8]. *Erysiphe alphitoides* complex has been described in Europe (Czechoslovakia, France, Germany, Lithuania, Netherlands, Poland, Romania, Russia, Serbia, Switzerland, and the United Kingdom) and Korea. However, the record of *E. alphitoides* on *Q. rubra* in Korea is erroneous because of the misidentification of the host plant. *Erysiphe extensa* var. *extensa* has been recorded in the United States (Missouri, New York, and Wisconsin) [3]. In conclusion, the *E. alphitoides* complex from Europe and *E. extensa* var. *extensa* from the United States were recorded for *Q. rubra*.

Recently, the *E. alphitoides* complex has been divided into several distinct species including *E. quercicola* [2,3,9]. Therefore, the identity of *E. alphitoides* on *Q. rubra* in European countries remains obscure. The observation of *E. quercicola* in Korea provides the first report of powdery mildew on *Q. rubra* in Asia. Two hypotheses can be put forward regarding this finding. One possibility is a biological invasion of the European isolate into East Asia, and another is the host expansion of the Korean isolate from the native *Quercus* spp. to the introduced *Q. rubra*. *Erysiphe quercicola* is one of the most common powdery mildew species on native *Quercus* spp. [10,11] and introduced *Q. robur* in Korea [12]. This is the first confirmed report on the identity of powdery mildew found on *Q. rubra* in Korea. This finding would be useful to understand the host expansion of *E. quercicola* and the biodiversity of powdery mildews on *Quercus* species.

CONFLICT OF INTERESTS

No conflict of interest was reported by the author(s).

ACKNOWLEDGMENTS

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, and Forestry (IPET), funded by the Ministry of Agriculture, Food, and Rural Affairs (MAFRA) (Project No. 320043-05).

REFERENCES

1. Shin HD, La YJ. Morphology of edge lines of chained immature conidia on conidiophores in powdery mildew fungi and their taxonomic significance. *Mycotaxon* 1993;66:445-51.
2. Takamatsu S, Braun U, Limkaisang S, Kom-Un S, Sato Y, Cunnington JH. Phylogeny and taxonomy of the oak powdery mildew *Erysiphe alphitoides* sensu lato. *Mycol Res* 2007;111:809-26.
3. Braun U, Cook RTA. Taxonomic manual of the Erysiphales (Powdery Mildews). CBS Biodiversity Series No. 11. CBS: Utrecht; 2012.
4. Bradshaw M, Tobin PC. Sequencing herbarium specimens of a common detrimental plant disease (powdery mildew). *Phytopathology* 2020;110:1248-54.
5. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol Biol Evol* 2013;30:772-80.
6. Vaidya G, Lohman DJ, Meier R. SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 2011;27:171-80.
7. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 2018;35:1547-9.
8. Farr DF, Rossman AY. Fungal databases [Internet]. Beltsville, MD: U.S. National Fungus Collections, ARS, USDA; 2022 [cited 2022 Feb 24]. Available from: <http://nt.ars-grin.gov/fungalatabases/>.
9. Takamatsu S, Ito H, Shiroya Y, Kiss L, Heluta V. First comprehensive phylogenetic analysis of the genus *Erysiphe* (Erysiphales, Erysiphaceae) I. The *Microsphaera* lineage. *Mycologia* 2015;107:475-89.
10. Lee HB, Kim CJ, Mum HY, Lee KH. First report of *Erysiphe quercicola* causing powdery mildew on ubame oak in Korea. *Plant Dis* 2011;95:77.
11. Lee HB, Nguyen TTT. First report of powdery mildew caused by *Erysiphe quercicola* on *Quercus mongolica* in Korea. *Plant Dis* 2017;101:1324.
12. Cho SE, Lee SH, Lee SK, Seo ST, Shin HD. First report of powdery mildew caused by *Erysiphe quercicola* on *Quercus robur* in Korea. *Plant Dis* 2018;102:1455.