

RESEARCH NOTE

First Report of *Sclerotinia sclerotiorum* Causing Sclerotinia Rot on *Ixeridium dentatum* in Korea

Myung Soo Park¹, Young Guk Kim², Sang Won Lee², Chun Geun Park², Yong Il Kim², Eun Song Lee², Jae Ki Chang², Tae Jin An^{2*}

¹School of Biological Sciences and Institute of Microbiology, Seoul National University, Seoul 08826, Korea

²National Institute of Horticultural and Herbal Science, Eumseong 27709, Korea

*Corresponding author: atj0083@korea.kr

Abstract

Sclerotinia rot was observed on *Ixeridium dentatum* cultivated as a succeeding crop in a garlic field in Seosan-si, Korea during the growing season in 2016 and 2017. Symptoms progressed from the initial irregular, water-soaked spots on main stems to wilting and eventually to plant death. White, cottony mycelia and black, irregular sclerotia formed on the basal stem and on soil surfaces. The optimal temperature of hyphal growth and sclerotia germination were 20°C and 25°C, respectively. Sequence analysis of the internal transcribed spacer (ITS) regions revealed that the three strains isolated from *Ixeridium dentatum* are grouped with *Sclerotinia sclerotiorum*. Three strains were identified as *Sclerotinia sclerotiorum* based on morphological features, ITS sequence, and pathogenicity test. To the best of our knowledge, this work is the first report of *Sclerotinia sclerotiorum* causing sclerotinia rot on *Ixeridium dentatum* in Korea.

Keywords: *Ixeridium dentatum*, New report, Sclerotinia rot, *Sclerotinia sclerotiorum*

OPEN ACCESS

Kor. J. Mycol. 2017 December, 45(4): 381-385
<https://doi.org/10.4489/KJM.20170045>

pISSN : 0253-651X
eISSN : 2383-5249

Received: 10 October, 2017

Revised: 30 October, 2017

Accepted: 3 November, 2017

© The Korean Society of Mycology



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ixeridium dentatum Tzvelev (family Asteraceae) is a perennial herb native to and widely distributed in Korea, Japan, and China [1]. The plant is well known for its anti-cancer, anti-oxidative and anti-allergenic activity [2, 3]. It is cultured in some areas, including Seosan, Dangjin, and Jinan. Sclerotinia rot symptoms were occasionally observed on *Ixeridium dentatum* cultivated as a succeeding crop in garlic fields in Seosan-si during the growing season of 2016 and 2017. Infected plants formed initially irregular, water-soaked spots on the base of the stems, wilted, then blighted and eventually died. Spherical or irregular sclerotia (2 to 7 mm in size) formed on the basal stem and on the soil surface (Fig. 1A, 1B). Cucumber mosaic virus (CMV disease), tomato spotted wilt virus (TSWV) and *Puccinia lactucae-debilis* (Rust) have been reported to be associated with diseases of *Ixeridium dentatum* in Korea [4]. The aim of the present study was to identify the causal

agent of sclerotinia rot on *Ixeridium dentatum*, based on morphological features, sequence analysis of the internal transcribed spacer (ITS) region and pathogenicity test.



Fig. 1. Symptom of sclerotinia rot on *Ixeridium dentatum* caused by *Sclerotinia sclerotiorum* and culture features. A, B, Infected plants; C, D, Symptoms induced by artificial inoculation; E, Mycelial mat and sclerotia produced on potato dextrose agar after 10 days at 20°C.

Sampling and Isolation

Infected tissues and sclerotia on the basal stem of *Ixeridium dentatum* were collected from several sites, where it was cultivated as succeeding crop in garlic fields at Seosan-si during 2016 and 2017. The tissues and sclerotia were surface-sterilized by dipping in 1% NaOCl solution for 1 min, rinsed three times with sterilized distilled water, and placed on water agar at 25°C for two days. The margins of each fungal hyphae growing from the tissue and sclerotia were transferred to potato dextrose agar (PDA; Difco, Detroit, MI, USA) plates. The isolated strains were maintained in a sterilized distilled water stock with 30% glycerol at -70°C for identification and pathogenicity tests in Herbal Crop Research Division (HCRD).

DNA extraction, PCR, sequencing, and phylogenetic analysis

Genomic DNA extraction and PCR amplification of the ITS region were performed using previously described methods [5]. DNA sequencing was performed at Macrogen (Seoul, Korea), using an ABI Prism 3700 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA). Sequences were assembled, proofread, edited using MEGA ver. 5.0 [6]. The ITS sequences of the present isolates were aligned with reference sequences downloaded from GenBank using the default settings of MAFFT v7 [7]. Neighbor-joining (NJ) trees were

constructed with MEGA 5 using Kimura 2-parameter model [8] and 1,000 bootstrap replicates. All the sequences were deposited in GenBank (accession nos. MF270175~MF270177). HCRD 16078, HCRD 16080, and HCRD 16084 isolated from infected tissues formed a monophyletic group with *Sclerotinia sclerotiorum* with 56% bootstrap value. The sequences of the three strains were identical to those of strains previously identified as *Sclerotinia sclerotiorum* (Fig. 2).




Fig. 2. Neighbor joining tree inferred from the internal transcribed spacer sequences of *Sclerotinia* species. Bootstrap values greater than 50 are presented at the nodes. The scale bar indicates the number of nucleotide substitutions per site. The isolates originated from *Ixeridium dentatum* are indicated in bold.

Culture and morphological characteristics

Pure cultures on PDA grew rapidly and formed aerial, white mycelia and turned gray to chocolate after 3 to 4 days. Fungal cultures formed spherical or irregular, dark-black sclerotia on the peripheral edges of the PDA, which ranged from 3 to 8 mm in size (Table 1). The optimal temperature for hyphal growth and sclerotia formation was 20°C, whereas that for sclerotia germination was 25°C. The morphological characteristics of the strains from sclerotinia rot of *Ixeridium dentatum* were similar to those of previously reported *S.*

sclerotiorum [9].

Table 1. Comparison between morphological features of the strains obtained from sclerotinia rot of *Ixeridium dentatum* and previously reported as *Sclerotinia sclerotiorum*

Characteristics		Present isolate	<i>Sclerotinia sclerotiorum</i> ^a
Colony	Color	white, gray to chocolate	White, grey to chocolate
	Size (mm)	2.5~6.0 × 2.6~8.5	3.0~10.0
Sclerotia	Shape	spherical or irregular	spherical or irregular
	Color	dark black	dark brown to black

^aDescribed by Kohn [9].

Pathogenicity test

Pathogenicity tests were conducted by placing five agar plugs (6 mm) of mycelium near the base of the stems of three healthy plants at soil line level. Non-inoculated plants were used as controls. All plants were kept in a dew chamber at 25°C and over 95% relative humidity. After three days, all inoculated plants showed symptoms, such as water soaked spots on the stem, covered with white mycelia, and then rotting, followed by wilting, blighting and eventually, death (Fig. 1C, 1D), whereas non-inoculated plants remained symptomless. *Sclerotinia sclerotiorum* was consistently re-isolated from the symptomatic tissue in full compliance with Koch's postulates.

Three Korean strains (HCRD 16078, HCRD 16080 and HCRD 16084) related to sclerotinia rot on *Ixeridium dentatum* were identified *S. sclerotiorum*, based on these morphological characteristics, sequence analysis, and pathogenicity test. This is the first report of *S. sclerotiorum* causing sclerotinia rot on *Ixeridium dentatum* in Korea.

Acknowledgements

This research was supported by the national joint agricultural research Program (Project No. PJ01187802) of the Rural Development Administration, Republic of Korea.

REFERENCES

1. Yook CS. Colored medicinal plants of Korea. Seoul: Academy Books; 1989.
2. Lee MR, Lee HY, Lee GH, Kim HK, Kim NY, Kim SH, Kim HR, Chae HJ. *Ixeris dentata* decreases ER stress and hepatic lipid accumulation through regulation of ApoB secretion. Am J Chin Med 2014;42:639-49.
3. Yi JM, Hong SH, Lee HJ, Won JH, Kim JM, Jeong DM, Baek SH, Lim JP, Kim HM. *Ixeris dentata* green sap inhibits both compound 48/80-induced anaphylaxis-like response and IgE-mediated anaphylactic response in murine model. Biol Pharm Bull 2002;25:5-9.
4. The Korean Society of Plant Pathology. List of plant diseases in Korea. 5th ed. Seoul:

- Korean Society of Plant Pathology; 2009.
5. Park MS, Fong JJ, Oh SY, Houbroken J, Sohn JH, Hong SB, Lim YW. *Penicillium jejuense* sp. nov., isolated from the marine environments of Jeju Island, Korea. *Mycologia* 2015;107:209-16.
 6. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011;28:2731-9.
 7. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 2013;30:772-80.
 8. Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 1980;16:111-20.
 9. Kohn LM. Monographic revision of the genus *Sclerotinia*. *Mycotaxon* 1979;9:365-444.