

RESEARCH ARTICLE

Lichen Genus *Porina* in VietnamSantosh Joshi¹, Dalip K. Upreti², Jae-Seoun Hur^{3,*}¹Department of Applied Science and Humanities, Invertis University, Bareilly (UP), India²Lichenology laboratory, CSIR-National Botanical Research Institute, Lucknow (UP), India³Korean Lichen Research Institute, Sunchon National University, Suncheon 57922, Korea

* Corresponding author: jshur1@sunchon.ac.kr

ABSTRACT


An identification key to twenty-nine species of *Porina* known from Vietnam is provided. In addition, new records of *Porina eminentior*, *P. meridionalis* and *P. nuculastrum* are described from the protected rain forests in southern Vietnam. A detailed taxonomic account of the newly reported species is provided and supported by its ecology, distribution, and illustrations.

Keywords: Corticolous, Nam Cat Tian National Park, Porinaceae, Taxonomy

INTRODUCTION

The cosmopolitan genus, *Porina* (Porinaceae: Ostropales), with more than 320 species worldwide, is most diverse in rather shaded habitats of tropical and subtropical regions [1-10]. The tropical climate of Vietnam is supported by prolonged humid conditions because of the large coastline surrounded by the South China Sea in the east and the Pacific Ocean in the south. These conditions are favorable to the growth of *Porina* on a range of substrates in tropical rainforests, seasonal forests, and wet lands in the country. The present study on this genus is a continuation of previous studies [11-16] and was conducted in Nam Cat Tien National Park (Fig. 1). The national park includes one of the largest areas of lowland tropical rainforests in southern Vietnam. Tree layer of the forest is comprised mainly of *Dipterocarpus alatus*, *D. intricatus*, *Dalbergia alata*, *D. mammosa*, *Azelia xylocarpa*, *Pterocarpus macrocarpus*, *Lagerstroemia calyculata*, *Tetrameles nudiflora*, *Anogeissus acuminata*, *Bambusa procera*, and *Gigantochloa* sp. Subtree layer is composed of *Actinodaphne pilosa*, *Beilschmiedia micranthopsis*, and *Illigera rhodantha*. *Amaranthus spinosus*, *Acanthus leucostachyus*, *Celosia argentea*, *Cyclacanthus coccineus* and *Ruellia repens* compose the shrub layer of the forest.

The checklist and preceding works on Vietnamese lichens reflect the great diversity of foliicolous species of *Porina* in the country (ca. 25 species), while a few were also reported on rock and bark [17,18]. Two species stated in the earlier account have now been accommodated in the genus *Strigula* as *S. phyllogena* (Müll. Arg.) R. C. Harris (= *Porina phyllogena* Müll. Arg.) and *S. platypoda* (Müll. Arg.) R. C. Harris (= *Porina platypoda* Müll. Arg.), whereas the previous report of *Porina consanguinea* Müll. Arg. from the country was uncertain in subsequent studies [17]; hence, it was not included in the current account.



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2019 September, 47(4): 303-11
<https://doi.org/10.4489/KJM.20190035>

Received: July 25, 2019
Revised: November 05, 2019
Accepted: November 08, 2019

© 2019 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The salient taxonomic features of *Porina* species are mostly shiny (corticate or ecorticate) thallus in different shades of greenish grey to olivaceous grey, containing algae either *Trentepohlia* (for bark and rock inhabiting species) or *Phycopeltis* (for leaf inhabiting species); perithecia present on thallus or immersed in thallus-dominated verrucae (Harada [6] proposed the term 'prominent thalloid exciple' replacing the 'thallus-dominated verrucae', applied previously to categorize the perithecial morphology by McCarthy [1]); pale brown to reddish brown or black, vestigial to well-developed involucrellum and hyaline, transversely three or more septate to muriform ascospores [1]. The genus is usually devoid of chemical compounds, but some species can give a K⁺ reddish or yellowish reaction on the thallus or on the fruiting bodies.

The present study was carried out in order to evaluate and enrich the diversity of Vietnamese lichens. The conducted work not only provides the noteworthy species of *Porina*, but also indicates the scope of non-lichenized fungi in unexplored areas of the country.




Fig. 1. Map of Vietnam showing the study area (Nam Cat Tien National Park).

MATERIALS AND METHODS

The interesting samples were selected from the lichen collection made by one of the authors (JSH) in December 2015 from the three provinces (Dong Nai, Lam Dong and Binh Phuoc) that circumscribed the national park. The field trip was organized by Dr. Nguyen Thi Thanh at Tay Nguyen University, Vietnam. The trip was conducted in the frame of internal joint program between Korea and Vietnam, supported by Korea National Research Foundation. The field studies did not involve any endangered or protected species. The material was preserved in the herbarium of the Korea National Arboretum, Korea (KH). The material was made available for taxonomic treatment in the Lichenology laboratory of CSIR-National Botanical Research Institute, Lucknow, India. The standard protocols were followed for identification. The morphological and anatomical characters were studied using a Leica S8APO stereo-zoom microscope and Leica DM500 compound microscope, respectively. Thin, hand-cut sections (10-15 sections) of perithecia, initially mounted in water, were studied for a range of structures and measurements. Cotton blue, 5% KOH and Lugol's iodine solution were used wherever required. Thin layer chromatography was performed in solvent system A according to the report by Orange et al. [19]. Illustrations were prepared using Corel Draw (ver. 12).

RESULTS AND DISCUSSION

Porina eminentior (Nyl.) P. M. McCarthy (Fig. 2A, 2B and 2C)

Lichenologist 32: 42 (2000).

Thallus epiperidermal, greenish grey to pale greenish grey, smooth to slightly verruculose, \pm glossy, continuous, sometimes cracked (due to bark texture), ≤ 150 μ m thick, corticate; algal layer *Trentepohlioid*, ≤ 100 μ m thick, medulla indistinct to endoperidermal; ascomata perithecioid, emergent, subglobose to hemispherical, 0.3-0.6 mm in diam.; ostiolar region pale brown to brown, 0.04-0.08 mm in diam.; involucrellum covered completely by a prominent thalloid exciple; proper exciple hyaline to pale brown, 15-20 μ m thick; centrum clear, 0.2-0.4 mm wide; asci 8-spored, $100-170 \times 25-35$ μ m; ascospores hyaline, broadly ellipsoidal, muriform, $40-70 \times 15-25$ μ m, multi-septate, perispore 3-5 μ m.

Chemistry: No lichen substances detected by thin layer chromatography.

Known distribution: In Vietnam, this species was found growing on tree twigs and most commonly associated with some sterile Porinaceae members in the wet tropical complex of the national park. *Porina eminentior* is also found in the Neotropics, East Africa, SE Asia, New Caledonia and the South Pacific [1].

Material examined: Vietnam, Dong Nai Province, Tan Phu district, Nam Cat Tien National Park, $11^{\circ}26'35''$ N $107^{\circ}24'19''$ E, 150 m elevation, on bark, 18 December 2015, Hur & Woo VN150213 (KH).

Notes: The species with muriform ascospores was assigned previously to the genus *Clathroporina* Müll. Arg. but later following the conservative concept outlined by Santesson [20] and McCarthy & Malcolm [21], Lücking & Vězda [22] included them in *Porina*. The Vietnamese specimen closely matches the original description of *Porina eminentior*, except that the examined sample has cracked rather than rimose thallus as described in McCarthy [1]. The thallus texture of our sample is somewhat similar to *P. nuculastrum*, but differs in ascospores with multi-separation.

***Porina meridionalis* P. M. McCarthy (Fig. 2D, 2E)**

Nova Hedwigia 58: 397 (1994).

Thallus epiperidermal, pale greenish grey, smooth, matt, continuous in large patches of ≤ 10 cm, flaking away from the bark in the middle part, delimited by brownish black prothallus, ecorticate, 50-70 μ m thick; algal layer *Trentepohlioid*, 20-40 μ m thick; medulla not apparent; ascomata perithecioid, semi-immersed to slightly emergent, hemispherical to subglobose, 0.1-0.3 mm in diam.; ostiole black, 0.01-0.03 mm in diam.; involucrellum dimidiate, covered mostly by a prominent thalloid exciple, 60-70 μ m thick; proper exciple hyaline to pale brown, 20-30 μ m thick; centrum clear, 0.25-0.27 mm thick; asci 8-spored, 90-130 \times 10-17 μ m; ascospores hyaline, elongate fusiform to subacicular, transversely 10-14-septate, 25-52 \times 5-8 μ m, perispore absent.

Chemistry: No lichen substances detected by thin layer chromatography.

Known distribution: This species appears to grow luxuriantly in large patches near coastal areas or mangroves in Vietnam and is collected from the thick and smooth trunk of trees in the protected forests of the park. The species was previously described in Tasmania [1].

Material examined: Vietnam, Dong Nai Province, Tan Phu district, Nam Cat Tien National Park, 11°27'27"N 107°22'11"E, 174 m elevation, on bark, 19 February 2015, Hur & Woo VN150389 (KH).

Notes: This species is distinct in producing elongate fusiform transversely 11-17(-21)-septate ascospores usually broader towards the distal end. McCarthy [1] reported the species with an inconspicuous ostiolar region, which is slightly conspicuous (0.01-0.03 mm in diam.) in this specimen. The Vietnamese sample is close to *Porina raphidiophora* (Nyl.) Müll. Arg., which differs mainly in the size of the perithecia and ascospores [1]. Although such minor variations may be acceptable within the broad concept of species delimitation, more collection is needed to observe the actual variations between the Vietnamese sample and the only report of this species from Tasmania, or more collection is prerequisite to describe the sample as noval.

***Porina nuculastrum* (Müll. Arg.) R. C. Harris (Fig. 2F, 2G)**

More Florida Lichens: 174 (1995).

Thallus epiperidermal, pale grey, smooth to verruculose due to emerging ascomata, glossy, continuous with black prothallus, <100 μ m thick; cortex 5-10 μ m thick; algal layer *Trentepohlioid*, 10-20 μ m thick; medulla white, crystalline, <60 μ m thick; ascomata perithecioid, scattered, distinctly emergent, convex to subglobose, 0.3-0.5 mm in diam., ostiole pale brown to dark brown, 0.1-0.3 mm; involucrellum apical to dimidiate, pale brown, covered completely by a prominent thalloid exciple; proper exciple hyaline to yellowish brown, 25-30 μ m thick; centrum clear, 0.3-0.5 mm wide; asci 8-spored, 200-250 \times 30-35 μ m; ascospores hyaline, narrowly to broadly fusiform, muriform (12-15 transverse septa and 1-3 longitudinal septa), 50-76 \times 12-20 μ m, perispore 2-3 μ m thick.

Chemistry: No lichen substances detected by thin layer chromatography.

Known distribution: This species was found growing in small and irregular patches on rough barked trees of the protected rainforests in Vietnam. The species has a common occurrence on tree bark as well

as on rocks. Worldwide, it is distributed in the Neotropics, Madagascar, the Philippines, Hong Kong, and neighboring countries [1].

Material examined: Vietnam, Dong Nai Province, Tan Phu district, Nam Cat Tien National Park, 11°26'35"N 107°24'19"E, 150 m elevation, on bark, 18 December 2015, Hur & Woo VN150198 (KH).

Notes: McCarthy [1] described this species with an inconspicuous ostiolar region. The examined material could be placed in *P. eminentior*, but multiple optical sections of perithecia showed very less septation in the measured ascospores. In contrast to the Australian specimens with an inconspicuous ostiole, the Vietnamese sample contains perithecia with a rather conspicuous ostiolar region. The ostiole conspicuity in the present sample may be the result of over maturity because only a single specimen from Vietnam was observed.




Fig. 2. New records from Vietnam. A, Habitus of *Porina eminentior*; B-C, Ascospores of *Porina eminentior*; D, Habitus of *P. meridionalis*; E, Asci and ascospores of *P. meridionalis*; F, Habitus of *Porina nuculastrum*; G, Ascospores of *Porina nuculastrum* (Scale bars: A, D, F = 1 mm; B, C, E, G = 50 µm).

Key to *Porina* species recorded from Vietnam

The taxonomic key characters of twenty-nine species are discussed. The previously known taxa were characterized following the descriptions provided by McCarthy [1], Upreti [19], Lücking & Vězda [22] and Pinokiyo & Singh [23] and presented below with some modifications.

1. Thallus foliicolous; algae *Trentepohlia* or *Phycopeltis* 2
- 1a. Thallus corticolous or saxicolous; algae *Trentepohlia* 23
2. Algae *Trentepohlia*; prothallus broad bluish black to grey black; thallus sterile or fertile 3
- 2a. Algae *Phycopeltis*; prothallus if present translucent or whitish; thallus fertile 4
3. Thallus sterile, isidiate; isidia abundant, cylindrical to coralloid *Porina distans*
- 3a. Thallus fertile; perithecia applanate, large, 0.6-1 mm in diam.; ascospores broadly fusiform, transversely 7(-9)-septate, $38-64 \times 6.5-10 \mu\text{m}$ *Porina imitatrix*
4. Perithecia immersed in thallus-dominated verrucae (thalloid exciple), mainly pale grey green, pale yellowish green or pale greyish brown 5
- 4a. Perithecia \pm superficial, not immersed in thallus dominated verrucae (thalloid exciple), yellowish white, in shades of brown or black 11
5. Ascospores 3-septate, $12-18 \times 2-3.5 \mu\text{m}$ ($13-25 \times 2-4 \mu\text{m}$ vide Lücking & Vězda, 1998); perithecia 0.15-0.2 mm in diam. *Porina albicera*
- 5a. Ascospores 7- or more septate 6
6. Thallus and perithecia furnished with dense cushions of soft hairs; perithecia convex to hemispherical; ascospores 7-15-septate, $45-74 \times 4-7 \mu\text{m}$ *Porina virescens*
- 6a. Thallus and perithecia glabrous; perithecia variously shaped; ascospores exclusively 7-septate (with a few exceptions), mostly less than $50 \mu\text{m}$ long 7
7. Perithecia subconical to conical; apex with a short flat-topped cylindrical extension; ascospores $28-46 \times 3-5 \mu\text{m}$ *Porina conica*
- 7a. Perithecia lens-shaped, hemispherical, or wart-shaped; if conical then apex lacking short flat-topped extension 8
8. Perithecia applanately lens-shaped; thallus with whitish short-stalked deeply concave disciform isidia, 0.1(-0.15) mm in diam. (0.4-0.7 mm in diam. vide Lücking & Vězda, 1998); ascospores $35-48 \times 3-4.5 \mu\text{m}$ [(35-)40-45 \times 3-4.5 μm vide Lücking & Vězda, 1998] *Porina mirabilis*
- 8a. Perithecia lens-shaped to hemispherical; thallus lacking isidia 9
9. Perithecia lens-shaped, 0.2-0.35(-0.4) mm in diam., with a blackish apical spot or a brownish apical cap; ascospores $23-33 \times 3-4 \mu\text{m}$ (27-35 \times 3-4.5 μm vide Lücking & Vězda 1998) *Porina atriceps*
- 9a. Perithecia hemispherical, concolorous or pale reddish brown at apices 10
10. Perithecia 0.2-0.3 mm in diam.; phycobiont cell rectangular, arranged in radiating rows; ascospores $21-27 \times 3 \mu\text{m}$ *Porina epiphylloides*
- 10a. Perithecia 0.25-0.4(-0.5) mm in diam.; phycobiont cells round to angular, irregularly arranged; ascospores $25-35 \times 3-4.5 \mu\text{m}$ *Porina epiphylla*

11. Perithecia very small, 0.08-0.1 mm in diam., yellowish white; ascospores bacillar, 1-septate, $10-12 \times 2-2.2 \mu\text{m}$ *Porina diaphana*
- 11a. Perithecia comparatively, brown to black, larger; ascospores 3- or more septate 12
12. Perithecia in shades of brown; involucrellum often containing algae 13
- 12a. Perithecia \pm black; involucrellum not or rarely containing algae 16
13. Perithecia finely and sparsely furnished with cushions of soft hairs, 0.2-0.26 mm in diam.; ascospores 7-septate, $25-33 \times 3-5 \mu\text{m}$ *Porina octomera*
- 13a. Perithecia not pilose; ascospores 3-septate 14
14. Perithecia subglobose, 0.17-0.26 mm in diam., constricted at the base; sides often with a slightly rough covering of thallus material; ascospores $17-26 \times 3-6 \mu\text{m}$ *Porina semecarpi*
- 14a. Perithecia convex to subconical 15
15. Perithecia 0.2-0.4 mm in diam.; ascospores $18-28 \times 2.5-4.5 \mu\text{m}$ *Porina limbulata*
- 15a. Perithecia 0.13-0.21 mm in diam.; ascospores $13-21 \times 2-3 \mu\text{m}$ *Porina rubentior*
16. Ascospores muriform, $85-110 \times 17-22 \mu\text{m}$; perithecia 0.35-0.50 mm in diam. *Porina foliicola*
- 16a. Ascospores transversely 3- or more septate 17
17. Ascospores 3-septate, $15-24 \times 3-4.5 \mu\text{m}$; perithecia applanate towards the margins, but usually with a conical to hemispherical centre, 0.2-0.41 mm in diam. *Porina chrysophora*
- 17a. Ascospores more than 3-septate 18
18. Perithecia convex to hemispherical; base usually spreading 19
- 18a. Perithecia subglobose to globose and attenuated at the base 22
19. Perithecia overgrown almost to the apex by a thin layer of thallus; ascospores 5(-7)-septate, $20-32 \times 4-5 \mu\text{m}$ *Porina corruscans*
- 19a. Perithecia not overgrown by the thallus; ascospores 5- or more septate 20
20. Ascospores 5-septate, $15-30 \times 5-7 \mu\text{m}$ *Porina subnitidula*
- 20a. Ascospores 7- or more septate 21
21. Ascospores 7-septate, $20-34 \mu\text{m}$ long *Porina cupreola*
- 21a. Ascospores 7-9-septate, $41-63 \mu\text{m}$ long (20-35 μm long vide McCarthy 2001) *Porina karnatakensis*
22. Perithecia 0.14-0.25 mm in diam.; sometimes greyish tomentose, ascospores 5-septate, $20-32 \times 3.5-6 \mu\text{m}$ *Porina nitidula*
- 22a. Perithecia 0.16-0.31 mm in diam.; ascospores mostly 7-septate, $22-42 \times 3.5-7 \mu\text{m}$ *Porina atrocoerulea*
23. Thallus saxicolous; perithecia hemispherical to subglobose, black, 0.23-0.48 mm in diam.; involucrellum uniformly black lacking algal cells; ascospores (3-)5-7(-9)-septate, $22-49 \times 3.5-6.5 \mu\text{m}$ *Porina guentheri*
- 23a. Thallus corticolous 24
24. Ascospores transversely septate 25
- 24a. Ascospores muriform 28
25. Ascospores elongate fusiform to subacicular, 10-14-septate, $25-52 \times 5-8 \mu\text{m}$; perithecia hemispherical to subglobose, 0.1-0.3 mm in diam. *Porina meridionalis*

25a. Ascospores fusiform	26
26. Ascospores (7-)9-11(-13)-septate	<i>Porina internigrans</i>
26a. Ascospores consistently 7-septate	27
27. Ascospores 4-9 μ m wide, perispore indistinct	<i>Porina tetracerae</i>
27a. Ascospores 6-13 μ m wide, perispore distinct	<i>Porina mastoidea</i>
28. Ascospores narrowly to broadly fusiform, 50-76 \times 12-20 μ m	<i>Porina nuculastrum</i>
28a. Ascospores broadly ellipsoidal, 40-70 \times 15-25 μ m	<i>Porina eminentior</i>

ACKNOWLEDGEMENTS

This paper was supported by Sunchon National University Research Fund in 2018 (Grant number: 2018-0287). SJ thanks Director, CSIR-National Botanical Research Institute, Lucknow for providing laboratory facilities and Invertis University, Bareilly (India) for research co-operation.

REFERENCES

- McCarthy PM. *Porina*. Fl Australia 2001;58A:106-52.
- Moon KH, Aptroot A. Pyrenocarpous lichens in Korea. *Bibl Lichenol* 2009;99:297-314.
- Cáceres MES, Dos Santos MWO, Mendonça CO, Mota DA, Aptroot A. New lichen species of the genera *Porina* and *Byssoloma* from an urban Atlantic rainforest patch in Sergipe, NE Brazil. *Lichenologist* 2013;45:379-82.
- Lücking R, Breuss O, Nelsen MP, Navarro E, Aptroot A, Chaves JL, Trest M, Will-Wolf S. *Porina squamulifera* (lichenized Ascomycota), a new species from tropical rainforest in Costa Rica with unique thallus morphology. *Herzogia* 2013;26:223-30.
- Tretiach M. *Porina pseudohibernica* sp. nov., an isidiate epiphytic lichen from central and south-eastern Europe. *Lichenologist* 2014;46:617-25.
- Harada H. Saxicolous and corticolous species of *Porina* (lichenized Ascomycota, *Porinaceae*) of Japan (part 1). *Lichenology* 2015;14:1-26.
- Orange A. A new freshwater *Porina* (*Porinaceae*, *Ostropales*) from Great Britain. *Lichenologist* 2015;26:223-30.
- Harada H. Saxicolous and corticolous species of *Porina* (lichenized Ascomycota, *Porinaceae*) of Japan (part 2). *Lichenology* 2015;14:1-26.
- Sobreira PNB, Aptroot A, Cáceres MES. Two new species of *Porina* (*Porinaceae*, lichenized Ascomycota) from submontane forest enclaves in Northeastern Brazil. *Phytotaxa* 2016;286:203-6.
- Upreti DK. Notes on corticolous and saxicolous species of *Porina* from India, with *Porina subhibernica* sp. nov. *Bryologist* 1994;97:73-9.
- Joshi S, Jayalal U, Oh SO, Nguyen TT, Dzong NA, Hur JS. A new species of *Graphis* and new lichen records from Vietnam, including a second worldwide report of *Sarcographina cyclospora*. *Mycobiology* 2014;42:17-21.
- Joshi S, Upreti DK, Oh SO, Nguyen TT, Nguyen AD, Hur JS. New records of crustose lichens and a lichenicolous *Arthonia* from Vietnam. *Mycotaxon* 2015;130:329-36.

13. Joshi S, Upreti DK, Nguyen TT, Nguyen AD, Oh SO, Hur JS. *Fissurina hemithecioides*, a new species and new records of *Graphidaceae* from Vietnam. *Cryptogamie Mycol* 2015;36: 383-97.
14. Joshi S, Upreti DK, Nguyen TT, Dunzg NA, Hur JS. New and interesting species in the family *Graphidaceae* (Ascomycota: *Ostropales*) from Vietnam. *Lichenologist* 2017;49: 259-68.
15. Joshi S, Upreti DK, Hur JS. Key to the lichen families *Pyrenulaceae* and *Trypetheliaceae* in Vietnam, with eight new records. *Mycotaxon* 2018;132:957-69.
16. Joshi S, Upreti DK, Lee BG. New records of *Arthoniaceae* from Vietnam. *Mycotaxon* 2018;133:103-12.
17. Aptroot A, Sparrius LB. Additions to the lichen flora of Vietnam, with an annotated checklist and bibliography. *Bryologist* 2006;109:358-71.
18. Nguyen TT, Joshi Y, Lücking R, Nguyen AD, Wang XY, Koh YJ, Hur JS. Seven new records of foliicolous lichens from Vietnam. *Mycotaxon* 2011;117:93-9.
19. Orange A, James PW, White FJ. Microchemical methods for the identification of lichens. London:British Lichen Society; 2001.
20. Santesson R. Follicolous lichens I. A revision of the taxonomy of the obligately foliicolous, lichenized fungi. *Symb Bot Ups* 1952;12:1-590.
21. McCarthy PM, Malcolm WA. The genera of trichotheliaceae. *Lichenologist* 1997;29:1-8.
22. Lücking R, Vězda A. Taxonomic studies in foliicolous species of the genus *Porina* (lichenized Ascomycotina: Trichotheliaceae) II. The *Porina epiphylla* group. *Willdenowia* 1998;28:181-225.
23. Pinokiyo A, Simha KP. Follicolous lichens of India. Dehradun: BSMPS; 2014.