

RESEARCH ARTICLE

Westerdykella reniformis: A New Record from Field Soils in Korea

Maresh Adhikari¹, Sang Woo Kim¹, Byeong Heon Gwon¹, Han Jun Ju¹, Hyang Burm Lee², and Youn Su Lee^{1,*}

¹Division of Biological Resource Sciences, Kangwon National University, Chuncheon 24341, Korea


²Division of Food Technology, Biotechnology & Agrochemistry, College of Agriculture and Life Sciences, Chonnam National University, Gwangju 61186, Korea

*Corresponding author: younslee@kangwon.ac.kr

ABSTRACT

During a survey of fungal diversity in different provinces of South Korea in 2017, a new fungal isolate was discovered. This fungal isolate was identified as *Westerdykella reniformis*, based on its morphological characteristics and phylogenetic analysis, using internal transcribed spacer (ITS) and 28S ribosomal DNA (28S rDNA) sequence data. To our knowledge, *W. reniformis* has not previously been reported in South Korea. Thus, in this study, we report a new record of a species from the *Dothideomycetes* class in Korea, and provide a detailed description with morphological illustrations.

Keywords: Diversity, ITS, *Westerdykella reniformis*, 28S rDNA



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2020 March, 48(1): 47-53
<https://doi.org/10.4489/KJM.20200005>

Received: February 27, 2020

Revised: March 20, 2020

Accepted: March 20, 2020

© 2020 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The class *Dothideomycetes* comprises the most diverse class within the Ascomycota, which is the largest known fungal phylum. They occur almost everywhere on earth and are found mainly as plant pathogens that cause serious crop losses [1]. *Torula thermophila* has been reclassified as *Scytalidium thermophilum*. *S. thermophilum* is characterized by intercalary and terminal conidial chains [2]. *Westerdykella* species are widespread saprobic fungi that have been isolated on a variety of substrates, such as: soil, mud, algal debris, dung, and plant material [3]. This genus is well-known by its cleistothecoid ascomata with small asci containing one-celled ascospores devoid of germ slits [4]. Knowledge of the taxonomy of some fungal genera like *Torula*, *Scytalidium* and *Westerdykella* in Korea is limited. Only one species of *Torula*, namely, *Torula caligans*, and one species of *Westerdykella*, namely, *Westerdykella dispersa* have been recorded in South Korea since 2006 [5].

A survey of Korean indigenous fungal diversity was conducted in Jeju and Ulsan, South Korea, in 2017. Fungal isolates showing different morphologies were isolated in soil samples from paddy fields and orchards. As a result, the previously unrecorded *Dothideomycetes*, *Westerdykella reniformis*, was discovered. This fungal species had not yet been reported in Korea. In this report, we present macro-morphological and micro-morphological characteristics of this new reported fungal isolate.

MATERIALS AND METHODS

Isolation of fungal isolates

Soil samples were collected in 2017 in different paddy fields located in Ulsan (35°31'45.36" N, 129°06'23.12" E) and Orchards located in Jeju (33°27'13.77" N, 126°19'51.23" E), South Korea. Soil samples were collected at a depth of 10-15 cm by removing crop debris. The soil samples were air dried and stored in sterile polythene bags at 4°C before use. Morphologically different fungal isolates were isolated on potato dextrose agar (PDA; Difco Laboratories, Detroit, Michigan, USA) supplemented with 100 µg L⁻¹ chloramphenicol using a conventional soil dilution technique [6]. The diluted soil suspensions were streaked on petri plates and incubated at 25°C for 5 days. Then, the morphologically different colonies that developed were aseptically transferred to fresh PDA plates to get pure cultures. The pure fungal isolates were finally kept on PDA slants at 4°C for further use.

Morphological examination

The macro-morphological characteristics of the fungal isolate KNU17-72 was determined on five different agar media, namely Malt extract agar (MEA), Potato dextrose agar (PDA), Yeast extract sucrose agar (YES), Czapek yeast extract agar (CYA) and Oatmeal agar (OA). Mycelial plugs of the fungal isolates were inoculated on petri plates of each medium and incubated at 25°C in the dark for 7 days. Colony characteristics were documented and a light microscope (Olympus BX50F-3 microscope, Olympus Optical Co., Ltd., Tokyo, Japan) was used to examine microscopic fungal structures. For micro-morphological examination, mounts of all the isolates that had grown on MEA were prepared and photomicrographs were taken using an Olympus BX50F-3 microscope (Olympus Optical Co., Ltd., Tokyo, Japan) equipped with a CMOS digital camera (KOPTIC Korea Optics, Seoul, Korea). Furthermore, the microscopic structures were characterized using scanning electron microscope (SEM) (LEO 1450VP Variable Pressure Scanning Electron Microscope, Carl Zeiss, Oberkochen, Germany).

Genomic DNA extraction, PCR, sequencing, and phylogenetic analysis

Genomic fungal DNA was isolated from mycelia of fungal isolates with the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany). The internal transcribed spacer (ITS) region of each sample was PCR amplified using primers ITS1 (5'-TCCGTAGGTGAACCTGCG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') [7]. In addition, the large subunit of 28S rDNA was amplified with universal primers LR0R (ACCCGCTGAACTTAAGC) [8] and LR7 (TCCTGAGGGAACTTCG) [9]. The PCR products were sequenced at Macrogen (Seoul, Korea) using an ABI PRISM 3730XL Analyzer (Life Technologies, Gaithersburg, MD, USA). The DNA sequences were subjected to BLAST analysis using GenBank database (Table 1). The phylogenetic tree was constructed by the neighbor-joining method using the Kimura's 2-parameter model [10] implemented in the program MEGA version 6 [11] with 1,000 bootstrap replicates.

The presently studied fungal isolate, KNU17-72, has been deposited in the culture collection site of the

National Institute of Biological Resources (NIBR, Incheon, Korea), under NIBRFG000500006. The nucleotide sequence of the ITS gene of the study isolate, KNU17-72, is deposited in the GenBank data base under accession number MH231762.

Table 1. The ITS and 28S rDNA gene sequences of *Westerdykella reniformis* used in this study, along with reference strains and their GenBank accession numbers.

No. Species	Isolate No.	GenBank accession No.	
		ITS	28S rDNA
1. <i>Preussia isomera</i>	CBS 388.78	GQ203763.1	GQ203723.1
2. <i>Preussia lignicola</i>	CBS 363.69	GQ203783.1	DQ384098.1
3. <i>Preussia australis</i>	S: Lundqvist 20884-a	GQ203773.1	GQ203732.1
4. <i>Preussia terricola</i>	CBS 317.65	NR 156524.1	GQ203725.1
5. <i>Preussia funiculata</i>	F: Huhndorf 2577	GQ203762.1	GQ203722.1
6. <i>Westerdykella centenaria</i>	BS 142400	NR 156002.1	NG_058468.1
7. <i>Westerdykella nigra</i>	CBS 416.72	GQ203800.1	GQ203800.1
8. <i>Westerdykella angulata</i>	CBS 610.74	NR 155956.1	NR_155956.1
9. <i>Westerdykella ornata</i>	CBS 379.55	NR 103587.1	NR_103587.1
10. <i>Westerdykella reniformis</i>	RKGE-35	JX235700.1	JX235704.1
11. <i>Westerdykella multispora</i>	CBS 383.69	GQ203799.1	GQ203799.1
12. <i>Westerdykella rapa-nuiensis</i> ined. GE-2013	CBS-604.97	JX235699.1	JX235703.1
13. <i>Westerdykella dispersa</i>	CBS 297.56 (T)	NR 111187.1	NR_111187.1
14. <i>Westerdykella aurantiaca</i>	FNBR-03	JN118571.1	JN118571.1

ITS: Internal transcribed spacer.

RESULTS

Morphological characterization

In this study, the morphologically different fungal isolate KNU17-72 was found in paddy field located in Jeju. Five different agar media, namely MEA, PDA, YES, CYA and OA were used to examine the macro-morphological characteristics of the isolate. Furthermore, the isolate were characterized according to their microscopic structures using compound and scanning electron microscopes. The macro and micro-morphological characteristics of the isolate are described below.

Morphological characteristics of the KNU17-72 isolate

Colony characteristics: The colony features of the KNU17-72 isolate on different agar media are presented in Fig. 1. Colonies on MEA reached a diameter of 25 mm after 7 days of incubation at 25°C. Colonies were faint brown with a velvety texture and reverse faint brown in color. Colonies on PDA reached approximately 25-27 mm in diameter, forming isabelline colonies with a velvety texture and reverse isabelline color. Colonies on YES attained a diameter of 27 mm after 7 days of incubation at 25°C. Colonies were khaki with a velvety texture, and reverse khaki color. Colonies on CYA attained a diameter of approximately 25-27 mm after 7 days of incubation at 25°C. Colonies were khaki in color, presented moderately velvety texture, and were reverse khaki. Colonies on OA grew slowly compared to other agar

media used, attaining a diameter of 17 mm after 7 days of incubation at 25°C. Colonies were isabelline with white boundaries, and a velvety texture, and were reverse dark grey in color.

Micromorphology: brownish black cleistothecia after three weeks; globose to sub-globose asci (12-15 × 16-18 µm); each ascus producing 32 ascospores. Ascospores had a distinct central groove; without pycnidial state.




Fig. 1. *Westerdykella reniformis* KNU17-72 grown on different media at 25°C for 7 days. (A-E) Colonies grown on malt extract agar (MEA), potato dextrose agar (PDA), yeast extract sucrose (YES), Czapek yeast extract (CYA), and oatmeal agar (OA) (top=obverse, bottom=reverse). (F) asci. (G) and (H) ascospores (scale bars: F=20 µm; G-H=10 µm).

Phylogenetic analysis

Molecular analysis was carried out by comparing the ITS sequences from KNU17-72 with other strains available in the GenBank database using the BLAST program. The ITS and 28S rDNA gene sequences of *Westerdykella reniformis* used in this study, along with reference strains and their GenBank accession numbers. The combined analysis of the internal transcribed spacer and 28S large subunit ribosomal RNA gene sequences revealed that the study isolate, KNU17-72, was *W. reniformis*, (Fig. 2). In addition, the combined ITS and 28S gene sequence analysis also revealed that the KNU17-72 isolate was phylogenetically most closely related to *W. reniformis* strain RKGE-35, existing along with *W. reniformis* RKGE-35 as a monophyletic clade (97% phenotypic similarity) in the phylogenetic tree. (Fig. 2).




Fig. 2. Neighbor-joining phylogenetic analysis using combined analysis of internal transcribed rDNA sequences and 28S large subunit ribosomal RNA gene sequences of *W. reniformis* KNU17-72. The species examined in this study are in bold. Bootstrap scores of $\geq 50\%$ are indicated at the nodes. The scale bar represents the number of substitutions per site. Sequences of type specimens are indicated by a symbol (T).

DISCUSSION

In this study, the fungal isolate KNU17-72, showing different morphologies, was isolated in 2017, from soil samples collected from different locations in Jeju, South Korea. The macro-morphology of the Korean isolate, *W. reniformis* KNU17-72, was very similar (Table 2) to the description given in a previous report by Ebead et al. [3]. Briefly, the colonies of our KNU17-72 isolate were velvety and faint brown to khaki in color. Notably, KNU17-72 grew slowly on OA agar medium compared to the other agar media used, reaching only 17 mm after 7 days of incubation. This result agrees with a previous report [3], which states that *W. reniformis* grows slowly on OA agar medium. In addition, KNU17-72 produced ascospores having central groves (Fig. 1H), which appears to be distinctive of *W. reniformis*, as has been reported

Table 2. Morphological characteristics of KNU17-72 and the reference species *W. reniformis* on different agar media at 25°C.

Characteristics	(KNU17-72), study isolate	<i>Westerdykella reniformis</i> , previous report ^a
Colony color	Colonies were faint brown, velvety texture	Velvety, faint brown
Colony diameter	Grown slowly on oatmeal agar (OA) compared to remaining media attaining 17 mm diam after 7 days of incubation	Slow growing attaining 16 mm diam on OA after 7 days of incubation
Cleistothecia	Brownish black cleistothecia after a month	Brownish black cleistothecia after 26 days
Asci	Globose to subglobose; produced 32 ascospores (12-15×16-18 μm)	Globose to subglobose, rarely ovoid (12-18×11-17 μm); produced 32 ascospores,
Ascospores	Black in color and kidney-shaped with concave a distinct central groove;	Black in color and kidney-shaped with concave a distinct central groove;
Pycnidial	Absent	Absent

^aFrom the description of Ebead et al. 2012 [3].

previously by Ebead et al. [3]. Globose to sub-globose asci, another distinctive feature of *W. reniformis*, have been noted on our isolate, KNU17-72 [3]. Furthermore, similar to other known *W. reniformis* isolates, pycnidia were not observed on our KNU17-72 isolate. Furthermore, the results of the combined analysis using the sequences of the internal transcribed spacer region and 28S large subunit ribosomal RNA gene sequences revealed that the KNU17-72 isolate is most closely related to *W. reniformis* strain RKGE-35, and exists along with *W. reniformis* RKGE-35 as a monophyletic clade (97% phenotypic similarity) in the phylogenetic tree. To the best of our knowledge, *W. reniformis* had not yet been reported in Korea and this is its first report.

ACKNOWLEDGEMENTS

This research was financed by the Ministry of Environment (MOE) of the Republic of Korea with the National Institute of Biological Resources (NIBR) grant (NIBR2014-01205) on survey and discovery of indigenous fungal species of Korea. This research study has also been sponsored by the University Industry Cooperation Foundation of Kangwon National University.

REFERENCES

1. Schoch CL, Crous PW, Groenewald JZ, Boehm EW, Burgess TI, De Gruyter J, De Hoog GS, Dixon LJ, Grube M, Gueidan C, et al. A class-wide phylogenetic assessment of Dothideomycetes. *Studies in mycology* 2009;1:64:1-5.
2. El-Sharouny H, Gherbawy YAMH, Abdel-Aziz F. Fungal diversity in brackish and saline lakes in Egypt. *Nova Hedwigia* 2009;89:437-50.
3. Ebead GA, Overy DP, Berru   F, Kerr RG. *Westerdykella reniformis* sp. nov., producing the antibiotic metabolites melinacidin IV and chetracin B. *IMA fungus* 2012;32:189-201.
4. Kruys A, Wedin M. Phylogenetic relationships and an assessment of traditionally used taxonomic characters in the Sporormiaceae (Pleosporales, Dothideomycetes, Ascomycota), utilizing multi-gene phylogenies. *Systematics and Biodiversity* 2009;7:465-78.
5. Lee YS, Jung HY, Lee HB, Kim SH, Shin KS, Eom AH, Kim C, Lee SY. National list of species of Korea. Ascomycota, Glomeromycota, Zygomycota, Myxomycota, Oomycota. Incheon: National Institute of Biological Resources; 2015.
6. Davet P, Rouxel F. Detection and isolation of soil fungi. Enfield (NH): Science Publishers; 2000.
7. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: A guide to methods and applications*. San Diego (CA): Academic Press; 1990. p. 315-22.
8. LoBuglio KF, Rogers SO, Wang CJK. Variation in ribosomal DNA among isolates of the mycorrhizal fungus *Cenococcum geophilum*. *Can J Bot* 1991;69:2331-43.
9. Vilgalys R, Hester M. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J Bacteriol* 1990;172:4238-46.
10. Kimura M. A simple method for estimating evolutionary rate of base substitution through

- comparative studies of nucleotide sequences. J Mol Evol 1980;16:111-20.
11. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol Biol Evol 2013;30:2725-9.