

RESEARCH NOTE

Damping-off of Edible Amaranth Caused by *Rhizoctonia solani* AG-4


Wan-Gyu Kim*, Hong-Sik Shim, Gyo-Bin Lee and Weon-Dae Cho
Global Agro-Consulting Corporation, Hwaseong 18330, Korea

*Corresponding author: wgkim5121@naver.com

ABSTRACT

Damping-off symptoms were frequently observed on young plants of edible amaranth (*Amaranthus mangostanus*) grown in vinyl greenhouses by farmers located in Goyang and Yangpyeong in Korea during disease surveys carried out in 2019 and 2020. The incidence of the disease was 1–20%. A total of eight isolates of *Rhizoctonia* sp. were obtained from the stems of the diseased plants. All the isolates were identified as *Rhizoctonia solani* AG-4, based on the morphological characteristics and anastomosis test. Three isolates of *R. solani* AG-4 were tested for pathogenicity on edible amaranth by artificial inoculation. All the tested isolates of *R. solani* AG-4 induced damping-off symptoms on the inoculated plants; these symptoms were similar to those observed on the vinyl greenhouse plants surveyed. This is the first report of *R. solani* AG-4 causing damping-off in edible amaranth.

Keywords: anastomosis group, damping-off, edible amaranth, pathogenicity, *Rhizoctonia solani*



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2020 September, 48(3): 325-328
<https://doi.org/10.4489/KJM.20200031>

Received: August 25, 2020

Revised: September 13, 2020

Accepted: September 16, 2020

© 2020 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Edible amaranth (*Amaranthus mangostanus* L.) is grown in the oriental countries as a vegetable crop. The crop is mostly cultivated in vinyl greenhouses in Korea. Damping-off symptoms were frequently observed on young plants of edible amaranth grown in vinyl greenhouses by farmers located in Goyang and Yangpyeong in Korea during disease surveys carried out in 2019 and 2020. The symptoms began on the soil surface parts of the plant stems. The infected parts of the stems receded and became thin, discolored and rotten. Diseased plants fell down and were wholly blighted (Fig. 1A and B). The incidence of the disease was 1–20% in the vinyl greenhouses investigated during the disease surveys (Table 1).

Rhizoctonia sp. was frequently isolated from the diseased plants of edible amaranth. A total of eight isolates of *Rhizoctonia* sp. were obtained from the stems of the diseased plants. All the isolates were identified as *Rhizoctonia solani* Kühn based on the morphological characteristics according to the descriptions of previous workers [1,2]. The isolates were tested to classify anastomosis groups using tester isolates of *R. solani* as previously conducted [3]. The tester isolates of *R. solani*, AG-1 (KACC 40101), AG-2-1 (KACC 40119), AG-2-2 (40125), AG-3 (KACC 40138), AG-4 (KACC 40139) and AG-5 (KACC 40146) were obtained from Agricultural Microbiology Division, National Institute of Agricultural Sciences

in Korea. All the tested isolates were classified as *R. solani* AG-4. Anastomosis reactions between an isolate and tester isolate of *R. solani* AG-4 were shown in Fig. 1C. The cultural appearance of *R. solani* AG-4 isolate is shown in Fig. 1D.




Fig. 1. Damping-off symptoms of edible amaranth and identification of the causal fungus, *Rhizoctonia solani* AG-4. A and B, damping-off symptoms on edible amaranth plants observed in the vinyl greenhouse; C, anastomosis reactions between an isolate and tester isolate of *R. solani* AG-4; D, a colony of *R. solani* AG-4 isolate grown on PDA at 25°C for 10 days; E, damping-off symptoms induced by artificial inoculation tests with *R. solani* AG-4 isolates. F, non-inoculated plants (control).

Table 1. Occurrence of damping-off of edible amaranth in vinyl greenhouses of farmers located in Goyang and Yangpyeong in Korea in 2019 and 2020.

Location investigated	Period investigated	No. of vinyl greenhouses investigated	No. of vinyl greenhouses with disease	% diseased plants ^a
Goyang	May, 2019	4	2	1-10
Yangpyeong	May, 2019	12	12	1-10
	April, 2020	16	16	1-20

^aOne hundred plants in each vinyl greenhouse were investigated with three replicates.

Three isolates of *R. solani* AG-4 were tested for pathogenicity on edible amaranth by artificial inoculation. Mycelial disks of 6 mm in diameter cut from the margins of actively growing cultures of each isolate on potato dextrose agar (PDA) were placed on stem bases of 25-day-old edible amaranth plants grown in circular plastic pots (9 cm in height, 10 cm in upper diameter and 7 cm in lower diameter) in the vinyl greenhouse. Inoculated plant pots were placed in plastic boxes (60×43×33 cm) with 100% relative humidity at room temperature (24-26°C) for three days. After that, the inoculated plant pots were taken out of the boxes and then kept indoors. Virulence of the isolates was rated based on the degree of damping-off symptoms induced at five days after inoculation. The inoculation test was performed with three replicates.

All the tested isolates of *R. solani* AG-4 induced damping-off symptoms on the inoculated plants (Fig. 1E). There was no symptom on the control plants (Fig. 1F). The symptoms on the plants induced by artificial inoculation were similar to those observed on the vinyl greenhouse plants surveyed. The isolates which induced symptoms on the plants were re-isolated from the symptoms.

R. solani was reported to cause damping-off in many crops [4-6]. Yang et al. [7] reported that *R. solani* AG-4HG-III causes stem canker and wirestem on green amaranth (*Amaranthus viridis* L.) and Chinese amaranth (*Amaranthus tricolor* L.). However, there has been no report on damping-off of edible amaranth caused by *R. solani*. This is the first report of *R. solani* AG-4 causing damping-off in edible amaranth.

ACKNOWLEDGEMENTS

This study was supported by a research grant (PJ014507012020) from Rural Development Administration, Korea.

REFERENCES

1. Parmeter JR Jr., Whitney HS. Taxonomy and nomenclature of the imperfect state. In: Parmeter JR Jr., editor. *Rhizoctonia solani*, Biology and Pathology. Los Angeles and London: University of California Press; 1970. p. 7-19.
2. Sneh B, Burpee L, Ogoshi A. Identification of *Rhizoctonia* species. The American Phytopathological Society, St. Paul, Minnesota, USA: APS Press; 1991.

3. Kim WG, Cho WD, Lee YH. Anastomosis groups and cultural characteristics of *Rhizoctonia solani* isolates from crops in Korea. Kor J Mycol 1994;22:309-24.
4. Kim WG. Pathogenicity of anastomosis groups and cultural types of *Rhizoctonia solani* isolates on crops. Korean J Plant Pathol 1996;12:21-32.
5. Kim WG, Cho WD, Ryu HY. Diagnosis and control of *Rhizoctonia* diseases on crops (in Korean). National Agricultural Science and Technology Institute, Suwon, Korea; 1995.
6. Farr DF, Rossman AY. Fungal Databases, U.S. National Fungus Collections, ARS, USDA; 2020 [cited 2020 August 14]. Available from: <https://nt.ars-grin.gov/fungaldatabases/>.
7. Yang GH, Chen HR., Naito S, Ogoshi A. Characterization of *Rhizoctonia solani* AG 4HG-III causing stem canker and wirestem on green amaranth and Chinese amaranth. J Phytopathol 2005;153:185-7.