

RESEARCH NOTE

Report on *Tuber huidongense*, a Truffle Species Previously Unrecorded in Korea

Hyeok Park¹, Ju-Hui Gwon¹, Jong-Chul Lee¹, Hyun Suk Kim², Deuk Sil Oh², and Ahn-Heum Eom^{1*}

¹Department of Biology Education, Korea National University of Education, Cheongju 28173, Korea


²Jeollanamdo Forest Resources Research Institute, Naju 58213, Korea

*Corresponding Author: eomah@knue.ac.kr

ABSTRACT

The fruiting bodies and roots of a *Tuber* species were collected from the rhizosphere of *Quercus dentata* in Pohang, Korea. The morphological characteristics of the ascoma and ectomycorrhizal roots were studied, and phylogenetic analyses were performed using the sequences of the internal transcribed spacer, large subunit rDNA, and β -tubulin DNA. Based on the features of the fruiting bodies, the species was identified to be *Tuber huidongense*, which has not been reported earlier in Korea. The morphological characteristics and phylogenetic analysis of *T. huidongense* are described in the present study.

Keywords: Ascomycota, Ectomycorrhiza, *Quercus dentata*



OPEN ACCESS

pISSN : 0253-651X

eISSN : 2383-5249

Kor. J. Mycol. 2020 December, 48(4): 505-510
<https://doi.org/10.4489/KJM.20200049>

Received: September 19, 2020

Revised: November 19, 2020

Accepted: November 30, 2020

© 2020 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fungi belonging to the genus *Tuber* F. H. Wigg. (Pezizales, Ascomycota) form a cleistothecial ascomata underground, which is commonly known as “Truffle” [1]. They have ectomycorrhizal (ECM) relationships with host plants [2]. *Tuber* species are widely distributed in the forests of the north temperate zone, including those in China and Japan [1]. Some of these have a particular aroma, owing to which they have high commercial value [1]. However, only a few reports have been published on *Tuber* spp. in Korea, which have primarily focused on the morphological characteristics of the ascoma [3] or the DNA sequences from the ECM roots without the fungi [4,5]. This is the first study to report the collection of the fruiting bodies and ECM root tips of *Tuber huidongense* Y. Wang in Korea, along with a description of its morphological characteristics and phylogenetic analyses.

The fruiting bodies were collected from the rhizosphere of *Quercus dentata* Thunb in Pohang, Korea (N35°57'16", E129°31'20"). The morphological characteristics of the fruiting bodies were evaluated, and genomic DNA was extracted from the ascoma using a DNeasy Plant Mini Kit (Qiagen GmVH, Hilden, Germany). For molecular identification, the ITS1F and ITS4 primers [6] were used to amplify the internal transcribed spacer (ITS) region, and the LR0R and LR16 primers [7] were used to amplify the large subunit (LSU) DNA. Nested PCR was performed to amplify the β -tubulin (TUB) DNA [8]. The first PCR experiment was performed using the Bt2a and Bt2b primers [9] for amplifying β -tubulin (TUB) DNA, in which the annealing temperature was set to 55°C. The first PCR product was diluted (1:10) and used as a template for the second PCR, which was performed using the *Tuber*-specific primers, Tubtubf and

Elytubr [8]. The annealing temperature was set to 63°C, and DNA sequence analysis was performed using the final round PCR products (SolGent, Daejeon, Korea). The sequences were determined using BLAST (available from the website of the National Center for Biological Information, NCBI), and the phylogenetic tree was constructed using the neighbor-joining method with MEGA7 [10]. The ECM roots of the oak tree were also collected, and their morphological characteristics were studied. The ECM root tip samples were identified based on the DNA sequences of the (ITS) region amplified using the ITS1F and ITS4 primers [6].

***Tuber huidongense* Y. Wang, Mycotaxon 83: 191 (2002) [MB#383644]**

Korean name: Huidong-deongi-beoseot (후이동덩이버섯); etymology from the species epithet derived from the name of the place where this species was first discovered

Classification: Ascomycota, Pezizales, Tuberaceae, *Tuber*

Morphology of ascoma: Cylindrical, short, elliptical, rugged, (10-)14(-20)×(8-)10(-12) mm in diam (Figs. 1A-C); Peridium: light or pale brown, approximately 140.97-153.15(-171.18) µm in thickness. Gleba: yellowish brown, paler than the peridium, white mycelia mixed in specific parts (Fig. 1D). Asci: hyaline, irregular, oval, or conical with rounded ends (Figs. 1E and 1F); 1-5 ascospores in each ascus, differing based on the number of ascospores; (42.13-)57.20(-77.71)×(28.87-)43.10(-57.32) µm Ascospore: light brown or yellowish brown; ellipsoid or fusiform; (16.95-)25.5(-33.47)×(14.19-)19.73(-25.85) µm (Fig. 1G). The reticular structure on the surface of the mature ascospore was similar to a turtle shell, with several spines present on the external surface; spines: (1.58-)3.06(-4.59) µm (Fig. 1H).

Specimen examined: Mt. Gwangjeongsan, Pohang-si, Gyeongsangbuk-do, Korea, N35°57'16", E129°31'20", August 3, 2020, *Tuber huidongense* GB20001, collected from the rhizosphere of *Quercus dentata*; GenBank accession No. MT940543 (ITS rDNA from Ascoma), MT940570 (LSU), MT950763 (TUB), and MW199065 (ITS rDNA from ECM root).



Fig. 1. Morphological characteristics of *Tuber huidongense*. ascomycete (A-C), peridium and gleba (D), asci (E, F), and ascospores (G, H) (scale bars: B, C=5000 µm, D=1000 µm, E=50 µm, F=20 µm, G, H=10 µm).

Phylogenetic analysis: BLAST results showed that the ITS sequence of GB20001 was closely related to that of *T. huidongense* FJ797884 (98.75% similarity), the LSU gene sequence was closely related to that of *T. huidongense* GU979099 (99.51% similarity), and the TUB gene sequence was closely related to that of *T. huidongense* GU979149.1 (99.41% similarity). The phylogenetic tree constructed using the ITS, LSU, and TUB sequences confirmed that the DNA sequences of GB20001 belonged to the same monophyletic group as those of *T. huidongense* (Fig. 2). Therefore, the phylogenetic results validated the identification of GB20001 as *T. huidongense*.




Fig. 2. Neighbor-joining phylogenetic tree of *T. huidongense* GB20001 ascoma based on concatenated alignment of internal transcribed spacer (ITS), large subunit (LSU) rDNA, and β -tubulin (TUB) DNA sequences. *Hypoxylon griseobrunneum* was used as an outgroup. Numbers on branches indicate bootstrap values (1,000 replicated).

Morphology of ECM roots: The ECM root tips were straight and cylindrical; their color varied from white to bright ivory; and white, fluffy emanating hyphae were observed on the surface (Figs. 3A-C). The cross section of the ECM showed the fungal mantle and Hartig net (Figs. 3D and 3E). The fungal mantle layer had a mosaic structure with irregularly shaped hyphae (Fig. 3F). The ECM root tips were identified as those of *T. huidongense* based on the DNA sequences of the ITS region (Fig. 4).




Fig. 3. Morphological characteristics of ectomycorrhizal colonized by *Tuber huidongense* from root of *Quercus dentata*. Mycorrhizal root tips (A-C); Transverse section of root tips stained with lactophenol cotton blue (D), outer mantle layer (E, F) (scale bars: A-C=500 μ m, D, E=100 μ m, F=20 μ m).




Fig. 4. Neighbor-joining phylogenetic tree of *T. huidongense* based on the alignment of internal transcribed spacer (ITS) rDNA sequences obtained from ectomycorrhizal root tips (MW199065) and ascoma of GB20001 (MT940543) in this study. *Hypoxylon griseobrunneum* was used as an outgroup. Numbers on branches indicate bootstrap values (1,000 replicates).

Wang and He [11] first reported the collection of *T. huidongense* in China; however, the morphological characterization and phylogeny were not clearly recorded, and there was no description of the ECM of *T. huidongense*. Deng et al. [12] reported additional morphological characteristics of *T. huidongense* and ectomycorrhiza formation with *Pinus armandii*. In this study, the ascoma and ascospores of GB20001 were found to be morphologically consistent with those of *T. huidongense*, which have been described in previous studies (Table 1). In addition, Deng et al. [12] reported that the ectomycorrhiza of *T. huidongense* with *P. armandii* had a bifurcate morphotype, while ectomycorrhiza with different host, *Q. dentata*, observed in this study, had an unbranched morphotype (Fig. 3). In conclusion, to our knowledge, this is the first report on *T. huidongense* in Korea.

Table 1. Morphological comparison of GB20001 with *Tuber huidongense* previously described.

Isolates	GB20001	<i>T. huidongense</i> [11]	<i>T. huidongense</i> [12]
Shape	Cylindrical, ellipsoid, rugged	Subglobose, ellipsoid	Subglobose, irregularly globose, ellipsoid
Size	(10-)14(-20)×(8-)10(-12) mm	5-25 mm in diam.	5-25 mm in diam.
Peridium	Pale brown to light brown, (140.97-)153.15(-171.18) µm thick	Yellowish-brown with red tint, 430-500 µm thick	Yellow-brown to red-brown, 150-300 µm thick
Asci	Hyaline, irregularly ellipsoid or coniform, 1-5 spored, (42.13-77.71)×(28.87-57.32) µm	(1)2-4 spored, Oblong(2-spored), subglobose to elliptical(3-spored), pyriform or ampulliform(4-spored), (45-60)×(37-50) µm	Ellipsoid or subglobose, 1-4(-5) spored, (42-70)×(35-55) µm
Ascospores	Light brown or yellowish brown, ellipsoid to fusiform, spiny-reticulate, (16.95-)25.57(-33.47)×(14.19-)19.73(-25.85) µm	Brown to pale brown, ellipsoid, spiny-reticulate, with bent points, (15-)23(-25)×(17-)18(-20) µm	Ellipsoid to narrowly ellipsoid, spiny-reticulate, yellowish brown, (22-45)×(16-29) µm
Spines	Straight, rarely curved, (1.58-)3.06(-4.59) µm high	Uncertain	Some curving at apex, (4-)6(-7) µm high

ACKNOWLEDGEMENT

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (IPET319106052HD050).

REFERENCES

1. Trappe JM. The orders, families, and genera of hypogeous Ascomycotina (truffles and their relatives). *Mycotaxon* 1979;9:297-340.
2. Pacioni G, Comandini O. *Tuber*. ectomycorrhizal fungi key genera in profile (Cairney WG, Chambers SM, eds), Berlin: Springer Verlag; 1999. p. 163.
3. Shin KS, Park JS, Yoshimi S. Note on *Tuber aestivum* subsp. *uncinatum* newly recorded in Korea. *Kor J Mycol* 1995;23:10-3.
4. Obase K, Lee JK, Lee SY, Chun KW. Diversity and community structure of ectomycorrhizal fungi in *Pinus thunbergii* coastal forests in the eastern region of Korea. *Mycoscience* 2011;52:383-91.

5. Obase K, Cha JY, Lee JK, Lee SY, Lee JH, Chun KW. Ectomycorrhizal fungal communities associated with *Pinus thunbergii* in the eastern coastal pine forests of Korea. *Mycorrhiza* 2009;20:39-49.
6. Gardes M, Bruns TD. ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Mol Ecol* 1993;2:113-8.
7. Moncalvo JM, Lutzoni FM, Rehner SA, Johnson J, Vilgalys R. Phylogenetic relationships of agaric fungi based on nuclear large subunit ribosomal DNA sequences. *Syst Biol* 2000;49:278-305.
8. Zampieri E, Mello A, Bonfante P, Murat C. PCR primers specific for the genus *Tuber* reveal the presence of several truffle species in a truffle-ground. *FEMS Microbiol Lett* 2009;297:67-72.
9. Glass NL, Donaldson GC. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol* 1995;61:1323-30.
10. Kumar S, Stecher G, Tamura K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 2016;33:1870-4.
11. Wang Y, He XY. *Tuber huidongense* sp. nov. from China. *Mycotaxon* 2002;83:191-4.
12. Deng XJ, Chen J, Yu FQ, Liu PG. Notes on *Tuber huidongense* (Tuberaceae, Ascomycota), an endemic species from China. *Mycotaxon* 2009;109:189-99.